



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ :C07D 217/24, C07C 39/38, 309/66, C07F
5/02, 5/04, 5/05

A1

(11) International Publication Number:

WO 96/15111

(43) International Publication Date:

23 May 1996 (23.05.96)

(21) International Application Number: PCT/US95/14896

(22) International Filing Date: 13 November 1995 (13.11.95)

(30) Priority Data:

08/339,958

15 November 1994 (15.11.94)

US

(71) Applicant: REGENTS OF THE UNIVERSITY OF MINNESOTA [US/US]; Morrill Hall, 100 Church Street S.E., Minneapolis, MN 55455 (US).

(72) Inventor: HOYE, Thomas, R.; 154 Montrose Place, St. Paul, MN 55104 (US).

(74) Agent: VIKSNINS, Ann, S.; Schwegman, Lundberg & Woessner, P.O. Box 2938, Minneapolis, MN 55402 (US).

(81) Designated States: AL, AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, LS, MW, SD, SZ, UG).

Published


With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: METHOD AND INTERMEDIATES FOR THE SYNTHESIS OF KORUPENSAMINES

(57) Abstract

A method of preparing a korupensamine or an analog thereof comprising: (a) reacting a compound of formula (III), wherein each of R¹ and R² is CH₃ or H, X is I, Y is (C₁-C₄)alkyl, benzyl or CHO, and each of R³ and R⁴ is (C₁-C₄)alkyl, benzyl, (C₂-C₅)acyl or an acid-labile hydroxy protecting group; with a compound of formula (IV), wherein R⁵ is benzyl, (C₂-C₅)acyl or an acid-labile hydroxy protecting group, R⁶ is B(OH)₂, and R⁷ is (C₁-C₄)alkyl; in the presence of a Pd(0) catalyst and an inorganic base in an organic solvent, to yield a compound of formula (V), wherein Y, R¹, R², R³, R⁴, R⁵ and R⁷ are as defined above for compounds of formula (III) and (IV). Additionally the intermediates of formula (III), wherein X is Br, Cl or I, Y is H, (C₁-C₄)alkyl, benzyl, or CHO, each of R¹ and R² is H or CH₃, R³ is a protecting group selected from the group consisting of (C₁-C₄)alkyl, benzyl, (C₂-C₅)acyl and an acid-labile hydroxy protecting group; and R⁴ is H or (C₂-C₅)acyl; or wherein X is Br, Cl or I, Y is H, (C₁-C₄)alkyl, benzyl, or CHO, each of R¹ and R² is H or CH₃, R³ is H or (C₂-C₅)acyl and R⁴ is a protecting group selected from the group consisting of (C₁-C₄)alkyl, benzyl, (C₂-C₅)acyl and an acid-labile hydroxy protecting group; and the intermediates of formula (IV), wherein R⁶ is Cl, Br, I, B(OH)₂, an anhydride or ester of B(OH)₂, or OSO₂R⁹, wherein R⁹ is (C₁-C₄)perfluoroalkyl, and each of R⁵ and R⁷ is H, (C₁-C₄)alkyl, benzyl, (C₂-C₅)acyl or an acid-labile hydroxy protecting group.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

METHOD AND INTERMEDIATES FOR THE
SYNTHESIS OF KORUPENSAMINES

Background of the Invention

5

Michellamines A (1), B (2), and C (3) constitute a family of anti-HIV, atropisomeric, naphthylisoquinoline alkaloids. All are fully protective against both HIV-1 and HIV-2 infected CEM-SS cells with EC₅₀ values of 2-13 μM. Michellamine B, the most studied and most prevalent of the group, completely protects MT-2 cells from both AZT-resistant and pyridone-resistant strains of HIV-1. The structure of michellamine B is shown below:



15

Michellamine B (2)


Michellamine B possesses a multilevel mode of action including an inhibition of the viral reverse transcriptase as well as blockage of cellular fusion and syncytium formation. In light of these promising properties, as well as favorable initial toxicity evaluation, michellamine B has been selected by the National Cancer Institute for IND-directed preclinical development. See, for example, K.P. Manfredi et al., *J.*

20

Med. Chem., **34**, 3402 (1991) and M.R. Boyd et al., J. Med. Chem., **37**, 1740 (1994).

The michellamines were isolated from a previously unidentified plant, *Ancistrocladus korupensis*--a liana found only in the rain forest of a limited region in Cameroon. Supply continuity and sufficiency are important concerns for further drug development. Atropisomers **1-3** are unique among known naphthylisoquinoline alkaloids in their dimeric nature, in the locus of the naphthalene to isoquinoline biaryl bond, and in the extent of free hydroxyl group adornment. The relative configurations of the stereogenic biaryl axes in each of **1-3** were established by identification of NOE interactions between the peri-H(1') and -H(1'') and one or the other of the diastereotopic protons at C(4) and C(4''). The absolute configurations at C(1)/C(1'') and C(3)/C(3'') were assigned by degradation to *R*-alanine and *R*-3-aminobutyric acid, respectively. See, M.R. Boyd et al., as cited above, G. Bringmann et al., Angew. Chem. Int. Ed. Eng., **32**, 1190 (1993) and G. Bringmann et al., Tetrahedron, **32**, 9643 (1994). The configurations at 5/8' and 8''/5''' of michellamines A, B and C are *S/S*, *S/R* and *R/R*, respectively.

Two syntheses of michellamine A were recently described by G. Bringmann et al., Tetrahedron, **32**, 9643 (1994), and T.R. Kelly et al., Tetrahedron Lett., **35**, 7621 (1994). An acyl derivative of a sample of the natural product korupensamine A (**4**), which co-exists with the michellamines in the plant, was oxidatively dimerized with silver oxide to yield a binaphthylidendione, which was reduced and deacylated to yield michellamine A. The structure of korupensamine A (**4**) and its atropisomer ("korupensamine C" (**4'**)) are shown below:


Korupensamine A (**4**)Korupensamine C (**4'**)

Compound 4' has been referred to as korupensamine B by G. Bringmann et al., *J. Org. Chem.*, **59**, 6349 (1994). In view of the ability to synthesize michellamines from these compounds in no more than five steps, a need exists for synthetic methods and intermediates which can be employed to prepare korupensamines.

Summary of the Invention

The present invention provides intermediates useful for the synthesis of korupensamines and thus for the synthesis of michellamines and their analogs.


For example, the present invention provides a compound of the general formula (I):



wherein X is Br, Cl or I; Y is H, (C₁-C₄)alkyl, benzyl, or CHO; each of R¹ and R² is H or CH₃ and each of R³ and R⁴ is H, (C₁-C₄)alkyl, benzyl, (C₂-C₅)acyl, or an acid-labile hydroxy protecting group such as (C₁-C₄)alkoxy(C₁-C₄)alkyl, tetrahydropyranyl, or (R⁸)₃Si, wherein each R⁸ is (C₁-C₄)alkyl. Preferably, X is I, R¹ and R² are CH₃, and R³, R⁴ and Y are the same protecting group, i.e., R³=R⁴=Y=benzyl.

As shown in compound 2, a broken line indicates a bond that extends below the plane of the ring, i.e., below the plane of the page, and a wedged line indicates a bond that extends above the plane of the page. Thus, to prepare korupensamines A and C, the 1*R*,3*R*-isomer of (I) is employed. However, the procedures disclosed by G. Bringmann et al., cited above, permit the preparation of all the 1,3-isomers of formula (I), wherein X=H; hence all the 1,3-isomers of formula (I) are considered to be within the scope of the invention.

The present invention also provides compounds of the formula (II):




(II)

wherein R^6 is Cl, Br, I, $B(OH)_2$, an anhydride or ester of $B(OH)_2$, or OSO_2R^9 ,
 wherein R^9 is (C_1-C_4) perfluoroalkyl, and each of R^5 and R^7 is H, (C_1-C_4) alkyl,
 5 benzyl, (C_2-C_5) acyl or an acid-labile hydroxy protecting group, as described above.
 Preferably, R^6 is Br or $B(OH)_2$, R^5 is an acid-labile protecting group, and R^7 is H or CH_3 .

The compound of formula II wherein R^6 is $B(OH)_2$ can be prepared
 from a compound of formula II wherein R^6 is halo, by lithiation and reaction of the
 10 lithiated compound with $B(OMe)_3$, following protection of the two OH groups, i.e.,
 wherein R^5 and R^7 are not H or acyl. The compound of formula II wherein R^6 is
 $B(OH)_2$ and R^5 and R^7 are not H or acyl can be coupled via Pd(0) catalyzed coupling
 with a compound of formula I, wherein X is I, $R^1=R^2=CH_3$, and R^3 , R^4 and Y are
 not H, to yield N- and 6,8,4',5' hydroxyl-protected korupensamines. Selective
 15 removal of the 5' hydroxyl protecting group, followed by oxidative 6'/6' coupling,
 reduction and, if necessary, removal of the remaining R^3 , R^4 and Y protecting
 groups, wherein R^7 is CH_3 , affords a mixture of michellamines A-C, which can be
 separated by chromatographic techniques.


Thus, a further aspect of the present invention is a method to prepare a
 20 korupensamine, preferably korupensamine A or B, or an analog thereof comprising:

(a) reacting a compound of the formula (III):




(III)

wherein each of R^1 and R^2 is CH_3 or H , X is I , Y is $(C_1-C_4)alkyl$, benzyl or CHO , and each of R^3 and R^4 is $(C_1-C_4)alkyl$, benzyl, $(C_2-C_5)acyl$ or an acid-labile hydroxy protecting group; with a compound of the formula (IV):



wherein R^5 is benzyl, $(C_2-C_5)acyl$ or an acid-labile hydroxy protecting group, R^6 is $B(OH)_2$, and R^7 is $(C_1-C_4)alkyl$; in the presence of a $Pd(0)$ catalyst and an inorganic base in an organic solvent, to yield a compound of the formula (V):



wherein Y , R^1 , R^2 , R^3 , R^4 , R^5 and R^7 are as defined above; and

(b) removing protecting groups R^3 , R^4 , R^5 and Y to yield a compound of formula V wherein R^1 and R^2 are each H or CH_3 , R^7 is $(C_1-C_4)alkyl$, and Y , R^2 , R^3 , R^4 , and R^5 are H . Preferably, R^1 , R^2 and R^7 are CH_3 , R^5 is an acid-labile protecting group, preferably methoxymethyl, that is subsequently removed by exposing V to dilute aqueous acid, and Y , R^3 and R^4 are benzyl that are subsequently removed by hydrogenolysis. Most preferably, the 1*R*,3*R*-isomer of III is employed, which yields a mixture of korupensamines A and C.


Brief Description of the Figures

Figure 1 is a reaction scheme summarizing the reaction of compounds 8 and 13 to yield korupensamine derivatives *S*-14 and *R*-14, and the conversion of these compounds to michellamines A-C.

Detailed Description of the Invention

As shown in Scheme I, following the general route developed by G. Bringmann et al., Angew. Chem. Int. Ed. Eng., **25**, 913 (1986) and G. Bringmann, et al., Liebigs Ann. Chem., 877 (1993), the non-racemic tetrahydroisoquinoline 7 (I, Y=H, R¹=R²=R³=R⁴=CH₃), was prepared from methyl 3,5-dimethoxybenzoate (5) via Raney nickel reduction of the non-racemic α -methylbenzylimine 6, following the methodology of D.E. Nichols et al., J. Med. Chem., **16**, 480 (1973). Demethylation of 7 with excess boron tribromide gave a diphenol amine•HBr salt, which was tribenzylated with benzyl bromide and cesium carbonate in DMF at room temperature (85%, two steps). Regiospecific iodination with iodine and silver sulfate gave C(5)-activated, benzyl protected 8 (66%), in accord with the methodology of W.W. Wy, Tetrahedron Lett., **34**, 6223 (1993). Likewise, direct bromination with Br₂ yields the corresponding brominated compound.

Scheme I.



5

Generally, demethylation of 7 yields a compound of formula I wherein Y, R^3 and R^4 are H, and the hydroxyl groups can be reacted with other protecting groups, such as those disclosed hereinabove. Likewise, the iodo moiety can be replaced by Br or Cl by a variety of halogen exchange reactions, such as by lithiation, following reaction with elemental halogen. Replacement of $(R)\text{-PhCH(Me)NH}_2$ with PhCH_2NH_2 and/or reaction with the corresponding 3,5-dimethoxyphenylacetaldehyde yields compounds of formula I wherein R^1 and/or R^2 is H. Synthesis of all the possible 1,3-isomers of 7, as well as compounds of formula 7 wherein one OMe group has been replaced by OH, is disclosed by G. Bringmann, et al., *Liebigs Ann. Chem.*, 877 (1993).

15


As shown in Scheme II, boronic acid 13 (II, $\text{R}^5=\text{CH}_2\text{OCH}_3$, $\text{R}^7=\text{CH}_3$, $\text{R}^6=\text{B(OH)}_2$) was efficiently prepared from methoxymethyl (MOM)-protected 2,4-dibromophenol (9) by a regioselective benzyne annulation reaction. Treatment of 9 with an excess of lithium cyclohexylisopropylamide and N,N-diethyl seneciarnide, as disclosed by M. Watanabe et al., *Chem. Pharm. Bull.*, 34, 2810 (1986), gave 12 (II, $\text{R}^5=\text{CH}_2\text{OCH}_3$, $\text{R}^7=\text{CH}_3$, $\text{R}^6=\text{Br}$) presumably by way of

20

benzyne **10** and lithium enolate **11**. Although the yield of this reaction was only 29%, the transformation was very reproducible. O-Methylation with methylsulfate, lithiation, and boronic acid synthesis with $B(OCH_3)_3$ followed standard protocols to yield **13**.

5 Reaction of a protected compound of formula II wherein $R^6=Li$ with Cl_2 or I_2 yields II, $R^6=Cl$ or I . Likewise, other acid-labile protecting groups can be used in place of $MeOCH_2$ in compound **9**, and $R^7=CH_3$ in formula II can readily be replaced with other protecting groups.

10

Scheme II.

As shown in Figure 1, the palladium(0) catalyzed cross-coupling of **8** with **13** provided an about 4:3 ratio of the hindered atropisomers *S*-**14** and *R*-**14** (40-80%). Palladium(0) catalyzed cross-coupling is typically carried out in the presence of a base and an organic solvent. A preferred embodiment of the invention utilizes tetrakis(triphenylphosphine)palladium(0) ($Pd(PPh_3)_4$) as the source of palladium(0) catalyst, saturated sodium bicarbonate ($NaHCO_3$) as the base and toluene as the organic solvent. Other useful sources of $Pd(0)$ catalysts include those disclosed in Larock et al. (U.S. Patent No. 5,233,059, Col. 6) and Blaser et al. (U.S. Patent No. 4,335,054, Col. 6 and Col. 7), which may alternatively be used in the method of present invention under conditions wherein $Pd(0)$ is generated. Bases useful in the present invention are those which are adequately soluble in the reaction medium. Although an inorganic base is preferred, an organic base can also be used.


Representative bases are disclosed at Col. 7 of the Larock et al. patent, and Col. 7 of the Blaser et al. patent, as cited above. Examples of suitable organic solvents include, in addition to the preferred toluene, tetrahydrofuran, ethers, glycol ethers, dimethylsulfoxide, dimethyl formamide, acetonitrile, acetamide, dimethylacetamide, and hexamethylphosphoramide.

Hydrolysis of the methoxymethyl (MOM) ethers of **14** gave the naphthols **15** (75-100%), which could be separated by careful normal-phase HPLC. Hydrogenolysis of the benzyl groups in a mixture of the naphthols **15** provided an about 4:3 mixture of korupensamine A and "C" atropisomers **4** and **4'**.

The mixture of tribenzylated naphthols **15** underwent remarkably efficient oxidative coupling with excess silver oxide in methylene chloride (or CDCl₃) at room temperature by the methodology of H. Laatsch, Liebigs Ann. Chem., 1321, (1980), to give the purple indigoids *R,S*-**16**, *S,S*-**16**, and *R,R*-**16** in an about 2:1:1 ratio (about 100%). The cross-ring quinones **16** could be reduced to the corresponding colorless binaphthols (sodium dithionite, H₂O, CH₂Cl₂ or NaBH₄, CH₂Cl₂, EtOH) and then perdebenzylated. More conveniently, direct exposure of **16** to one atmosphere of hydrogen in methylene chloride/methanol over 10% Pd/C resulted in simultaneous reductive bleaching of the indigoid and complete hydrogenolysis of the six benzyl groups. Michellamines A-C were cleanly (as judged from the crude ¹H NMR spectrum) produced with nearly quantitative mass recovery. Separation of a small portion on amino-bonded phase [7:1 CH₂Cl₂:0.1 weight % (NH₄)₂CO₃ in methanol] has thus far provided a pure sample of michellamine A (**1**) along with an about 2:1 mixture of michellamines B (**2**) and C (**3**), as determined by NMR analysis.

The invention will further be described by reference to the following detailed examples.


Example 1. Preparation of 1(*R*), 3(*R*)-1,2,3,4-tetrahydro-6,8-dihydroxy-1,3-dimethylisoquinoline hydrobromide salt (101)



10

(1*R*, 3*R*)-1,2,3,4-tetrahydro-6,8-dimethoxy-1,3-dimethylisoquinoline was prepared as described in G. Bringmann et al., Liebigs Ann. Chem., 877 (1993) and 50.7 mg (0.2 mmol) placed in an oven dried flask (5 mL r.b.) containing CH₂Cl₂ (1 mL) and a magnetic stir bar. The flask was sealed with a rubber septum and the atmosphere was exchanged for nitrogen. The reaction mixture was cooled to -78 °C and a BBr₃ solution (1 mL, 4.3 equiv, 1 M in CH₂Cl₂) was added via syringe. The reaction mixture was immediately allowed to warm to room temperature and stir. After 10 h, the flask was cooled to -78 °C and carefully quenched with 1.5 mL of MeOH. The stir bar was removed and the reaction mixture was concentrated in vacuo to yield a brown oil. MeOH (3.5 mL) was added to dissolve the oil and the reaction mixture was concentrated again. This quenching procedure was repeated 6-8 times until the hydrobromide salt **101** (62.8 mg, 100%) was isolated as brown crystals; ¹H NMR (500 MHz, CD₃OD): δ 6.23 [d, J = 1.8 Hz, Ar-H(7)], 6.12 [d, J = 2.1 Hz, Ar-H(5)], 4.64 [q, J = 6.7 Hz, CHCH₃], 3.75 [ddq, J = 11.6, 4.6 and 6.5 Hz, CH₂CHCH₃], 2.98 [dd, J = 17.4 and 4.6 Hz, CH_{ax}H_{eq}CHCH₃], 2.75 [dd, J = 17.4 and 11.6 Hz, CH_{ax}H_{eq}CHCH₃], 1.59 [d, J = 7.0 Hz, CHCH₃], and 1.46 [d, J = 6.4 Hz, CH₂CHCH₃]; ¹³C NMR (125 MHz, CD₃OD): δ 158.97, 156.10, 133.65, 112.61, 107.01, 101.94, 49.35, 45.35, 34.59, 19.23, and 18.33; m.p. (range): 140-143 °; Anal. Calcd for C₁₁H₁₆NO₂Br: C, 48.19; H, 5.88. Found: C, 48.35; H, 5.69.

**Example 2. Preparation of Tribenzylprotected Tetrahydroisoquinoline
(102)**




5

To a stirred solution of 1(R), 3(R)-1,2,3,4-tetrahydro-6,8-dihydroxy-1,3-dimethyl-isoquinoline hydrobromide salt (0.39 g, 1.4 mmol) in 15 mL of dry DMF was added benzylbromide (1.70 g, 10.0 mmol), followed by the addition of cesium carbonate (2.40 g, 7.4 mmol). After being stirred for 6 h at room temperature, the reaction mixture was poured into H₂O (100 mL), and EtOAc (100 mL) was added. The organic layer was washed with brine, dried over Na₂SO₄, and concentrated in vacuo. The residue was purified by flash chromatography (hexanes/EtOAc/Et₃N; 9:1:0.1) to yield tribenzyl-protected tetrahydroisoquinoline **102** (0.57 g, 86 %) as a thick yellow oil.; ¹H NMR (500 MHz, CDCl₃): δ 7.42-7.21 [m, benzyl ArH], 6.42 [d, J = 2.0 Hz, ArH(7)], 6.34 [d, J = 2.0 Hz, ArH(5)], 4.99 [s, O(6)CH₂Ph], 4.98 [d, J = 12.0 Hz, O(8)CH₂H_bPh], 4.95 [d, J = 12.0 Hz, O(8)CH_aH_bPh], 4.01 [q, J = 7.0 Hz, ArCHCH₃], 3.82 [d, J = 14.0 Hz, NCH₂H_bPh], 3.52 [ddq, J = 10.5, 4.5 and 6.5 Hz, CH₂CH_bCH], 3.32 [d, J = 14.0 Hz, NCH_aH_bPh], 2.63 [dd, J = 17.0 and 10.5 Hz, CH_aCH_bCH], 2.58 [dd, J = 17.0 and 4.5 Hz, CH_aCH_bCH], 1.34 [d, J = 6.5 Hz, CH₃(1)], and 1.26 [d, J = 6.5 Hz, CH₃(3)]; ¹³C NMR (75 MHz, CDCl₃): δ 157.6, 157.1, 137.1, 136.7, 129.0, 128.7, 128.5 [5C], 128.4 [2C], 128.3, 128.0 [2C], 127.9, 127.6 [2C], 126.9 [2C], 126.4, 105.5, 98.3, 70.0, 69.6, 51.2, 50.0, 45.7, 32.6, 19.9, and 19.5; IR (neat NaCl plates): 2967, 1603, 1454, and 1149 cm⁻¹; Anal. calcd for C₃₂H₃₃NO₂: C, 82.90; H, 7.17. Found: C, 82.89; H, 6.95.

25

Example 3. Preparation of Iodide (8)




5

A solution of compound **102** (0.48 g, 1.0 mmol) in 10 mL of EtOH and 2 mL of CH₂Cl₂ was added slowly to a stirred mixture of iodine (0.53 g, 2.1 mmol) and silver sulfate (0.69 g, 2.2 mmol) in 10 mL of EtOH. After being stirred at room temperature for 16 h, the yellow solid was removed by filtration and the filtrate was concentrated in vacuo. The resulting residue was dissolved in CH₂Cl₂ (100 mL). This solution was washed with saturated NaHCO₃, H₂O, dried over Na₂SO₄, and concentrated in vacuo. The residue was purified by flash chromatography (hexanes/EtOAc; 9:1) to yield iodide **8** (0.40 g, 66 %) as a thick oil; ¹H NMR (500 MHz, CDCl₃): δ 7.49-7.18 [m, benzyl ArH], 6.41 [s, ArH(7)], 5.07 [s, O(6)CH₂Ph], 4.98 [d, J = 12.0 Hz, O(8)CH₂H_bPh], 4.94 [d, J = 12.0 Hz, O(8)CH₂H_aPh], 4.01 [q, J = 6.5 Hz, CHCH₃], 3.82 [d, J = 14.0 Hz, NCH₂H_bPh], 3.51 [ddq, J = 12.0, 4.0 and 6.5 Hz, CH_aH_bCHCH₃], 3.20 [d, J = 14.0 Hz, NCH₂H_aPh], 2.66 [dd, J = 17.5 and 4.0 Hz, CH₂H_bCH], 2.42 [dd, J = 17.5 and 12.0 Hz, CH₂CH_aCH], 1.34 [d, J = 6.5 Hz, CH₃(1)], and 1.31 [d, J = 6.5 Hz, CH₃(3)]; ¹³C NMR (75 MHz, CDCl₃): δ 157.4, 156.1, 141.3, 139.5, 137.1 [2C], 128.9 [7C], 128.5 [2C], 128.2, 127.4 [2C], 127.2 [3C], 126.9, 124.3, 97.7, 71.6, 70.3, 51.9, 50.1, 46.9, 39.3, 20.2, and 20.1; IR (neat NaCl plates): 2971, 1585, 1324, and 1062 cm⁻¹; Anal. calcd for C₃₂H₃₂INO₂: C, 65.20; H, 5.47. Found: C, 65.39; H, 5.73.

25

Example 4. Preparation of 5-Bromo-(1*R*, 3*R*)-1,2,3,4-tetrahydro-6,8-dimethoxy-1,3-dimethylisoquinoline (103)



5


To a solution of tetrahydroisoquinoline **7** (50.0 mg, 0.23 mmol) in CH_2Cl_2 (0.5 ml) was added Br_2 (13 μL , 0.25 mmol). After stirring for 10 min the reaction was diluted with Et_2O , washed successively with with saturated K_2CO_3 , saturated $\text{Na}_2\text{S}_2\text{O}_3$, and brine. The organics were then dried over 4 A molecular sieves, filtered, and concentrated in vacuo. The resulting residue was triturated with Et_2O and a white solid was filtered (18.4 mg, 27% of brominated tetrahydroisoquinoline **103** as the HBr salt) The remaining material was purified by MPLC (hexanes/ EtOAc ; 1:1 with 3% Et_3N) to yield an additional quantity (39.4 mg, 57%) of the brominated species (84% total); ^1H NMR (200 MHz, CDCl_3): δ 9.6 [vds, 1H, NH], 6.39 [s, Ar-H(7)], 4.84 [q, $J = 6.7$ Hz, CHCH_3], 3.88 [s, OMe], 3.85 [s, OMe], 3.73 [ddq, $J = 11.6, 4.5$, and 6.2 Hz, CH_2CHCH_3], 3.19 [dd, $J = 17.7$ and 4.5 Hz, $\text{CHaH}_b\text{CHCH}_3$], 2.92 [dd, $J = 17.7$ and 11.6 Hz, $\text{CHaH}_b\text{CHCH}_3$], 1.81 [d, $J = 6.2$ Hz, CH_2CHCH_3], and 1.70 [d, $J = 6.7$ Hz, CHCH_3]; LRMS (EI): m/z 298 ($\text{M}^+ - 1$, <1), 286 (97), 284 (100), 269, 256, 226, 204, 190, 176, 162, 147, 131, 103, 91, 77, 51, and 42 (all <5).

15

20


Example 5. Preparation of *N*-benzyl-(1*R*, 3*R*)-1,2,3,4-tetrahydro-6,8-dimethoxy-1,3-dimethylisoquinoline (104)

25



Into a stirred solution of (1*R*, 3*R*)-1,2,3,4-tetrahydro-6,8-dimethoxy-1,3-dimethyl-isoquinoline (114 mg, 0.5 mmol) and benzyl chloride (137 mg, 1.1 mmol) in methyl ethyl ketone was added K₂CO₃ (320 mg, 2.3 mmol). The resulting mixture was heated to reflux for 24 h, after which time it was cooled down and poured into H₂O. Et₂O was added and the organic layer was washed with H₂O, brine, dried over MgSO₄, and concentrated in vacuo. The crude product was purified by MPLC (hexanes/EtOAc/Et₃N; 9:1:0.3) to yield N-benzyl-(1*R*, 3*R*)-1,2,3,4-tetrahydro-6,8-dimethoxy-1,3-dimethyl-isoquinoline (148 mg, 93 %) as a colorless oil; ¹H NMR (500 MHz, CDCl₃): δ 7.40-7.20 [m, benzyl ArH], 6.28 [d, J = 2.0 Hz, ArH(7)], 6.23 [d, J = 2.0 Hz, ArH(5)], 3.66 [q, J = 6.5 Hz, ArCHCH₃], 3.82 [d, J = 14.0 Hz, NCH₂H_bPh], 3.78 [s, O(6)CH₃], 3.70 [s, O(8)CH₃], 3.50 [ddq, J = 10.5, 5.0 and 6.5 Hz, CH₂H_bCHCH₃], 3.29 [d, J = 14.0 Hz, NCH₂H_bPh], 2.63 [dd, J = 17.0 and 10.5 Hz, CH₂H_bCH], 2.58 [dd, J = 17.0 and 5.0 Hz, CH₂H_bCH], 1.30 [d, J = 6.5 Hz, CH₃(1)], and 1.25 [d, J = 6.5 Hz, CH₃(3)]; ¹³C NMR (75 MHz, CDCl₃): δ 158.5, 158.4, 141.7, 136.8, 128.4 [2C], 128.1 [3C], 126.4, 104.1, 96.4, 55.2, 55.1, 51.4, 49.8, 45.8, 32.6, 20.0, and 19.5; LRMS (EI): *m/z* 296 (M⁺-CH₃, 100), and 91 (44); IR (neat NaCl plates): 2965, 1605, and 1148 cm⁻¹; Anal. calcd for C₂₀H₂₅NO₂: C, 77.14; H, 8.09. Found: C, 77.30; H, 8.06.


Example 6. Preparation of N-benzyl-(1*R*, 3*R*)-1,2,3,4-tetrahydro-6,8-dihydroxy-1,3-dimethylisoquinoline hydrobromide salt (105)



Into a 15 mL culture tube was placed N-benzyl-(1*R*, 3*R*)-1,2,3,4-tetrahydro-6,8-dimethoxy-1,3-dimethylisoquinoline (62 mg, 0.2 mmol) dissolved in acetic acid (1 mL). To this solution were added sodium iodide (120 mg, 0.8 mmol) and concentrated aqueous hydrobromic acid (49%, 2 mL). The mixture was heated at 100 °C for 3 hours and then cooled to 0 °C, at which time light yellow crystals precipitated out of solution. Vacuum filtration with a glass fritted Buchner funnel gave

the N-benzyl-(1R, 3R)-1,2,3,4-tetrahydro-6,8-dihydroxy-1,3-dimethyl-isoquinoline hydrobromide salt [105] (48 mg, 66%) as light yellow crystals .

5 **Example 7. Preparation of N-benzyl-(1R,3R)-5-bromo-1,2,3,4-tetrahydro-6,8-dimethoxy-1,3-dimethylisoquinoline (106)**




10

Into a 25 mL round bottom flask was placed N-benzyl-(1R, 3R)-1,2,3,4-tetrahydro-6,8-dimethoxy-1,3-dimethylisoquinoline (288 mg, 0.9 mmol) dissolved in methylene chloride (3 mL). Bromine (156 mg, 1.0 mmol) in methylene chloride (1 mL) was added to the solution. The mixture was stirred for 3 h at room temperature and then diluted with methylene chloride (20mL) and washed with H₂O (2 x 5 ml). The organic layer was dried with sodium sulfate and concentrated in vacuo to yield 360 mg of crude material. Separation on MPLC (hexane/ethyl acetate; 9:1, 3% Et₃N) gave N-benzyl-(1R, 3R)-5-bromo-1,2,3,4-tetrahydro-6,8-dimethoxy-1,3-dimethylisoquinoline [106] (282 mg, 78%) as light yellow oil; ¹H-NMR (CDCl₃, 200 MHz): δ 7.31 [m, 5H, Ph], 6.4 [s, Ar-H(7)], 3.97 [q, J = 6.6 Hz, CHCH₃], 3.91 [s, OMe] 3.84, [d, J = 14.4 Hz, CH_cH_dPh], 3.76 [s, OMe], 3.52 [ddq, J = 11.6, 6.0, and 4.6 Hz, CH₂CHCH₃], 3.19 [d, J = 14.4 Hz, CH_cH_dPh], 2.74 [dd, J = 17.7 and 4.6 Hz, CH_aH_bCHCH₃], 2.45 [dd, J = 17.7 and 11.6 Hz, CH_aH_bCHCH₃], 1.35 [d, J = 6.0 Hz, CH₂CHCH₃], and 1.31 [d, J = 6.6 Hz, CHCH₃]; ¹³C-NMR (CDCl₃, 200 MHz): δ 157.0, 154.4, 140.9, 136.1, 128.3 [2C], 128.0 [2C], 126.3, 122.1, 104.5, 94.3, 56.3, 55.3 [2C], 51.3, 49.4, 45.7, 33.1, and 19.8; LRMS (EI): m/z 374 (M⁺-15, 70), 360, 294, 268, 226, 203, 190, 162, 145 (all <5), 91 (100), 65, and 39.


30

Example 8. Preparation of Iodide (107)



5 A solution of N-benzyl-(1R, 3R)-1,2,3,4-tetrahydro-6,8-dimethoxy-
 1,3-dimethyl-isoquinoline (198 mg, 0.61 mmol) in 5 mL of EtOH was added slowly
 to a stirred mixture of iodine (333 mg, 1.3 mmol) and silver sulfate (468 mg, 1.5
 mmol) in 10mL of EtOH. After being stirred at room temperature overnight, the
 yellow solid was removed by filtration and the filtrate was concentrated in vacuo. The
 10 resulting residue was dissolved in 40 mL of CH₂Cl₂. The solution was washed with
 saturated NaHCO₃, H₂O, dried over Na₂SO₄, and concentrated in vacuo. The crude
 product was purified by flash chromatography (hexanes/EtOAc; 9:1) to yield iodide
107 (200 mg, 75 %) as a white solid; ¹H NMR (500 MHz, CDCl₃): δ 7.35-7.20
 [m, 5H, Ph], 6.36 [s, ArH(7)], 3.90 [q, J = 6.5 Hz, ArCH₂CH₃], 3.89 [s,
 15 O(6)CH₃], 3.81 [d, J = 14.5 Hz, NCH₂H_bPh], 3.75 [s, O(8)CH₃], 3.49 [ddq, J =
 11.5, 4.5 and 6.5 Hz, CH₂H_bCHCH₃], 3.16 [d, J = 14.5 Hz, NCH₂H_bPh], 2.63
 [dd, J = 17.5 and 4.5 Hz, CH₂H_bCH], 2.39 [dd, J = 17.5 and 11.5 Hz, CH₂H_bCH],
 1.33 [d, J = 6.5 Hz, CH₃(1)], and 1.29 [d, J = 6.5 Hz, CH₃(3)]; ¹³C NMR (75
 20 MHz, CDCl₃): δ 158.5, 156.8, 141.1, 139.1, 128.4 [2C], 128.1 [3C], 126.4,
 123.2, 94.0, 56.6, 55.3, 51.7, 49.6, 46.6, 38.7, 19.9, and 19.8; IR (neat NaCl
 plates): 2966, 1586, 1453, 1326, 1207, and 1072 cm⁻¹.

Example 9. Preparation of 2,4-Dibromo-1-methoxymethoxybenzene (9)




Into a 500-mL round bottom flask equipped with a soxlet and a condensor were placed 2,4-dibromophenol (32.0 g, 0.13 mole), dimethoxymethane (200 mL, 2.26 mole), p-toluenesulfonic acid monohydrate (2.24 g, 12.0 mmol) and CH₂Cl₂ (200 mL). The soxlet extractor was filled with 3 A and 4 A molecular sieves.

- 5 The reaction mixture was heated to reflux for 24 h after which time the soxlet extractor was filled with freshly activated sieves. The reaction mixture was heated to reflux for another 24 h. After this period of time, Et₃N (10 mL) was added. The reaction mixture was stirred for 5 min, and concentrated to dryness. The residue was dissolved in CH₂Cl₂ (400 mL) and the resulting solution was washed with 5 %
10 NaOH (400 mL), H₂O (400 mL), dried over Na₂SO₄, and concentrated in vacuo. The crude product was purified by flash chromatography (hexanes/EtOAc; 6:1) to yield 2,4-Dibromo-1-methoxymethoxybenzene [9] (32.9 g, 88 %) as a light yellow oil; ¹H NMR (CDCl₃, 300 MHz) δ 7.67 [s, ArH(3)], 7.33 [d, J = 8.7 Hz, ArH(5)], 7.02 [d, J = 8.7 Hz, ArH(6)], 5.21 [s, OCH₂OCH₃], and 3.49 [s, OCH₃]; LRMS
15 (EI) *m/z* 298 (M⁺, 3), 296 (M⁺, 5), 294 (M⁺, 3), and 45 (100).

Example 10. Preparation of Naphthol (12)


20



- To a stirred solution of isopropylcyclohexylamine (7.5 mL, 0.45 mmol) in 60 mL of THF at -78 °C under N₂ was added *n*-BuLi (20.0 mL, 50.0 mmol, 2.5 M in hexanes). The mixture was stirred for 20 min, warmed to 0 °C, and then stirred for 1 h. The mixture was cooled to -78 °C and solution of N,N-diethyl-3,3-dimethylacrylamide (2.10 g, 13.0 mmol) in 40 mL of THF was added. This mixture was stirred at -78 °C for 1 h. The cold bath was removed and the reaction mixture was allowed to warm to -20 °C over a period of 10 min. A solution of 2,4-
- 25

dibromo-1-methoxymethoxybenzene (9) in 30 mL of THF was added. The reaction mixture was stirred overnight at room temperature, and then quenched with saturated NH_4Cl . Et_2O was added and the solution was washed with H_2O , brine, dried over Na_2SO_4 , and concentrated in vacuo. The crude product was purified by flash chromatography (hexanes/ EtOAc ; 9:1) to yield naphthol 12 (1.15 g, 29 %) as a brown oil; ^1H NMR (CDCl_3 , 300 MHz): δ 9.31 [s, OH], 7.54 [d, $J = 8.4$ Hz, ArH(6)], 7.49 [s, ArH(4)], 6.82 [s, ArH(2)], 6.81 [d, $J = 8.4$ Hz, ArH(7)], 5.46 [s, OCH_2OCH_3], 3.55 [s, OCH_3], and 2.47 [s, $\text{ArCH}_3(3)$]; ^{13}C NMR (CDCl_3 , 75 MHz): δ 154.3, 153.5, 139.5, 134.5, 129.6, 118.2, 115.5, 114.5, 113.6, 107.3, 95.9, 56.9, and 22.0; LRMS (EI): m/z 298 (M^+ , 13), 296 (M^+ , 11), 128 (5), 115 (9), and 45 (100).


Example 11. Preparation of Bromide (111)




To a stirred solution of dimethyl sulfate (2.49 g, 20.0 mmol) in 20 mL of CH_2Cl_2 was added a solution of Bu_4NBr (2.19 g, 6.8 mmol) and NaOH (0.50 g, 12.0 mmol) in 15 mL of H_2O and a solution of naphthol 12 (1.07 g, 3.6 mmol) in 10 mL of CH_2Cl_2 . The reaction mixture was stirred at room temperature for 18 h. The organic and aqueous layers were separated and the aqueous layer was extracted with 20 mL of CH_2Cl_2 . The combined organic layers were washed with H_2O , dried over Na_2SO_4 , and concentrated in vacuo. The crude product was purified by MPLC (hexanes/ EtOAc ; 9:1) to yield 111 (0.83 g, 74 %) of as a white solid; ^1H NMR (CDCl_3 , 300 MHz): δ 7.63 [s, ArH(5)], 7.62 [d, $J = 8.1$ Hz, ArH(3)], 6.86 [d, $J = 8.1$ Hz, ArH(2)], 6.75 [s, ArH(7)], 5.22 [s, OCH_2OCH_3], 3.94 [s, $\text{ArO}(8)\text{CH}_3$], 3.58 [s, OCH_2OCH_3], and 2.51 [s, $\text{ArCH}_3(6)$]; ^{13}C NMR (CDCl_3 , 75 MHz): δ 156.8, 154.0, 137.8, 135.0, 130.4, 119.5, 118.1, 115.3, 112.9, 109.3, 96.8, 56.5,

56.4, and 22.2; LRMS (EI): m/z 312 (M^+ , 18), 310 (M^+ , 19), 282 (15), 280 (16), 231 (2), 128 (14), and 45(100).

5 **Example 12. Preparation of Compound (113)**




15 In a 250 mL round bottomed flask a solution of
lithiumisopropylcyclohexylamide was prepared from isopropylcyclohexylamine (6.21
g, 7.23 mL, 44 mmol) in THF (50 mL) and butyllithium (2.5 M in hexane, 17.6 mL,
44 mmol). The solution was cooled to - 78 °C under nitrogen and N,N-diethyl-3,3-
20 dimethylacrylamide [109] (3.10 g, 20 mmol) in THF (20 mL) was added. After
stirring for 30 min at - 78 °C, the solution was warmed to room temperature and
stirred for 5 h. The reaction mixture was then cooled to - 78 °C, and 1,4-
dibromoanisole [112] (5.852 g, 22 mmol) in THF (30 mL) was added via syringe.
The solution was stirred at 0 °C for 24 h and then quenched with saturated aqueous
25 ammonium chloride (100 mL) and diluted with ether (30 mL). The organic layer was
separated and the aqueous layer was extracted with ether (3 x 40 mL). The
combined organic layers were washed with brine (40 mL), dried with sodium sulfate
and concentrated in vacuo to give 4.60 g of crude material. Flash chromatography of
the crude material (hexane/ethyl acetate; 9:1) yielded product 113 (1.17 g, 22%) as a
30 light yellow oil .

Example 13. Preparation of Compound (114)

10 A solution of dimethyl sulfate (2.52 g, 20.0 mmol) in methylene chloride (20 mL) was prepared in a 100 mL round bottom flask. To the flask was added a solution of tetrabutyl-ammonium bromide (2.25 g, 7.0 mmol) and sodium hydroxide (400 mg, 10.0 mmol) in water (15 mL). To this mixture was added a solution of compound **113** (1.335 g, 5 mmol) in methylene chloride (10 mL). The reaction mixture was stirred at room temperature for 18 h and then diluted with methylene chloride (20 mL). The organic and aqueous layers were separated, and the aqueous layer was extracted with of methylene chloride (20 mL). The combined organic layers were washed with water (10 mL), dried with sodium sulfate, and concentrated in vacuo. The resulting residue was purified by flash chromatography (hexane/ethyl acetate; 9:1) to obtain compound **114** (1.32 g, 94%) as a white solid .

15

20

Example 14. Preparation of Compound (116)

30


In a 100 mL round bottom flask, 7-methyl-Juglone [**115**] (242.4 mg, 1.2 mmol) was dissolved in chloroform (30 mL). Water (15 mL) and sodium

dithionite (627 mg, 3.6 mmol) were added to this and the mixture was stirred at room temperature for 1 h. When TLC analysis showed no starting material remaining, the organic layer was separated, washed with brine, dried with sodium sulfate and concentrated in vacuo to yield compound 116 (245 mg, 100%) as a white solid.

5

Example 15. Preparation of Compound (117)

10




15

Compound 116 (204 mg, 1.0 mmol) was placed in a 50 mL round bottom flask with methylene chloride (10 mL) and pyridine (0.32 mL, 4.0 mmol). The mixture was cooled to - 5 °C and triflic anhydride (0.20 mL, 1.15 mmol) was added. The reaction mixture was stirred at room temperature for 12 h. Methylene chloride (20 mL) was added and the mixture was washed with water, dried with sodium sulfate and concentrated in vacuo. Purification by flash chromatography (Hexanes/ethyl acetate; 12:1) yielded product 117 (182 mg, 54%) as a white solid.

20

25


Example 16. Preparation of Compound (118)

30

Into a 15 mL round bottom flask was placed monotriflate **117** (50 mg, .
0.15 mmol) dissolved in methylene chloride (2 mL). To this solution was added
acetic anhydride (0.04 mL, 0.45 mmol) and pyridine (0.072 mL, 0.90 mmol). The
reaction mixture was stirred at room temperature for 9 h. When TLC analysis showed
5 no remaining starting material, the mixture was diluted with methylene chloride (10
mL), washed with water, dried with sodium sulfate and concentrated in vacuo to give
compound **118** (56 mg, 100%) as a white solid.


Example 17. Preparation of Boronic Acid (13)

10



15

To a stirred solution of **111** (0.83 g, 2.7 mmol) in 30 mL of THF at
-78 °C under N₂ was added *n*-BuLi (1.3 mL, 3.2 mmol, 2.5 M in hexanes). The
resulting mixture was stirred for 15 min and then cannulated into a solution of
20 B(OMe)₃ (0.65 mL, 5.7 mmol) in 30 mL of THF. The reaction mixture was stirred
for another 15 min, then warmed to room temperature and stirred for 2 h. The
reaction mixture was quenched with 10 % HCl, diluted with Et₂O, washed with H₂O,
brine, dried over MgSO₄, and concentrated in vacuo to yield boronic acid **13** (0.74 g,
100 %) as a brown solid. This compound was used without further purification. The
25 structure of boronic acid [**13**] was confirmed by derivatization to boronate ester **119**;
GC: t_R = 13.3 min; column: DB-5, 6 m x 0.1 mm x 0.1 μm film; temp prom: 50 °C /
2 min / 20 °C min⁻¹ / 250 °C / 10 min; LRMS (EI): *m/z* 344 (M⁺, 100), 314 (46),
300 (20), 270 (17), and 45 (75).

Example 18. Preparation of Boronic Anhydride (120)

5

Into a 25 mL flame dried flask was placed bromide **111** (1.10 g, 3.54 mmol) freshly distilled THF (10 mL). Under nitrogen, the solution was cooled to -78 °C and then n-butyllithium (2.5 M in hexane, 1.7 mL, 4.2 mmol) was added. The resulting solution was stirred for 15 minutes, after which time a precipitate appeared.

Trimethyl borate (1.7 mL, 14.4 mmol) was added to the flask and a clear solution formed. The mixture was stirred at -78 °C for 30 min and then at room temperature for 2 h. The mixture was quenched with saturated aqueous ammonium chloride (10 mL), concentrated, and diluted with of methylene chloride (20 mL). The organic and aqueous layers were separated and the aqueous layer was neutralized with 10% aqueous hydrochloric acid. The aqueous layer was extracted with methylene chloride (2 x 20 mL), and the combined organics were washed with H₂O (10 mL) and dried over sodium sulfate. Concentration in vacuo yielded 1.0 g of crude material as a caramel colored residue. Precipitation from methylene chloride by addition of hexanes gave the anhydride **120** (548 mg, 60%) as a white solid; ¹H-NMR (CDCl₃, 300 MHz): δ 8.72 [s, ArH(1)], 8.52 [s, J = 7.7 Hz, ArH(7)], 7.08 [d, J = 7.8 Hz, ArH(6)], 6.76 [s, ArH(3)], 5.37 [s, OCH₂OCH₃], 3.97 [s, ArOCH₃], 3.62 [s, OCH₂OCH₃], and 2.43 [s, ArCH₃].

25

Example 19. General Procedure for the Palladium(0)-Mediated Biaryl Coupling Reactions


30

To a stirred solution of aryl iodide in toluene (0.05 M) was added 2 equivalents of boronic acid (or its derivatives). Saturated NaHCO₃ (1/2 volume of toluene) was then added, followed by the addition of 20 mol % of Pd(PPh₃)₄. The reaction mixture was sealed under N₂ in a culture tube and heated to 110 °C for 20 h.

- 5 After this period of time, EtOAc and brine were added. The organic layer was extracted, dried over Na₂SO₄, and concentrated in vacuo. The crude product was purified by silica gel chromatography.

A. Preparation of Compound (14)

10




15

- Compound **14** was obtained from boronic acid **13** and iodide **8** in 81% yield with a 4:3 ratio of **S-14** to **R-14**. The product was purified by MPLC (hexanes/EtOAc/Et₃N; 3:1:0.1); ¹H NMR of **S-14** (500 MHz, CDCl₃): δ 7.39-6.90 [m, ArH(6' and 7') and benzyl ArH], 6.77 [s, ArH(1')], 6.69 [s, ArH(3')], 6.53 [s, ArH(7)], 5.31 [s, OCH₂OCH₃], 5.02 [s, O(6)CH₂Ph], 4.87 [d, J = 12.5 Hz, O(8)CH₂H_bPh], 4.81 [d, J = 12.5 Hz, O(8)CH_aH_bPh], 4.12 [q, J = 6.5 Hz, PhCHCH₃], 3.98 [s, O(4')CH₃], 3.72 [d, J = 14.5 Hz, NCH₂H_bPh], 3.65 [s, OCH₂OCH₃], 3.37 [ddq, J = 11.5, 4.0, and 6.5 Hz, CH_aH_bCHCH₃], 3.30 [d, J = 14.5 Hz, NCH_aH_bPh], 2.36 [s, CH₃(2')], 2.22 [dd, J = 17.5 and 4.0 Hz, CH₂H_bCH], 2.00 [dd, J = 17.5 and 11.5 Hz, CH_aH_bCH], 1.41 [d, J = 6.5 Hz, CH₃(1)], and 1.01 [d, J = 6.5 Hz, CH₃(3)]; ¹H NMR of **R-14** (500 MHz, CDCl₃): δ 7.39-6.90 [m, ArH(6' and 7') and benzyl ArH], 6.86 [s, ArH(1')], 6.70 [s, ArH(3')], 6.51 [s, ArH(7)], 5.31 [s, OCH₂OCH₃], 5.03 [d, J = 12.0 Hz, O(6)CH₂H_bPh], 4.97 [d, J = 12.0 Hz, O(6)CH_aH_bPh], 4.86 [d, J = 12.5 Hz, O(8)CH₂H_bPh], 4.81 [d, J = 12.5 Hz, O(8)CH_aH_bPh], 4.11 [q, J = 6.5 Hz,
- 20
- 25
- 30

PhCHCH₃], 3.98 [s, O(4')CH₃], 3.77 [d, J = 14.5 Hz, NCH₂H_bPh], 3.65 [s, OCH₂OCH₃], 3.37 [ddq, J = 14.0, 4.0, and 6.5 Hz, CH₂H_bCHCH₃], 3.35 [d, J = 14.0 Hz, NCH₂H_bPh], 2.36 [s, CH₃(2')], 2.25 [dd, J = 17.0 and 14.0 Hz, CH₂H_bCH], 1.92 [dd, J = 17.0 and 4.0 Hz, CH₂H_bCH], 1.39 [d, J = 6.5 Hz, CH₃(1)], and 1.05 [d, J = 6.5 Hz, CH₃(3)]; IR (neat NaCl plates): 2967, 1584, 1052, and 733 cm⁻¹.

B. Preparation of Compound (121)




Into a 15 mL culture tube were placed N-benzyl-(1R, 3R)-5-iodo-1,2,3,4-tetrahydro-6,8-dimethoxy-1,3-dimethylisoquinoline [107] (66 mg, 0.15 mmol) and toluene (3 mL). To this solution was added compound 120 (58 mg, 0.23 mmol), which resulted in the formation of a slurry. A minimum amount of ethanol was added to change the slurry to a clear solution. To the resulting solution was added tetrakis(triphenylphosphine)palladium(0) (17 mg, 0.015 mmol) and saturated aqueous sodium bicarbonate (1.5 mL). The atmosphere was exchanged for N₂ and the reaction mixture was heated at 110 °C for 12 hours. When TLC showed no substrate boronate left, the organic and aqueous layers were separated and the aqueous layer was extracted with ethyl acetate (2 x 5 mL). The combined organics were washed with brine, dried over sodium sulfate and concentrated in vacuo. The resulting residue was purified by silica gel chromatography (hexane/ethyl acetate ; 9:1, 3% Et₃N) to yield a mixture of compounds 121a and 121b (50 mg, 60%) as a white solid; ¹H NMR of 121a (500 MHz, CDCl₃): δ 7.39-7.21 [m, benzyl ArH], 7.17 [d, J = 8.0 Hz, ArH(7')], 7.07 [d, J = 7.5 Hz, ArH(6')], 6.74 [s, ArH(1')], 6.68 [s, ArH(3')], 6.49 [s, ArH(7)], 5.31 [s, OCH₂OCH₃], 4.00 [q, J = 6.5 Hz, NCHCH₃], 3.97 [s, O(4')CH₃], 3.85 [s, O(6)CH₃], 3.72 [d, J = 14.5, NCH₂H_bPh], 3.65 [s, O(8)CH₃], 3.64 [s, OCH₂OCH₃], 3.36 [ddq, J = 11.0, 6.5 and 4.0 Hz, CH₂H_bCHCH₃], 3.26 [d, J = 14.0 Hz, NCH₂H_bPh], 2.35 [s, CH₃(2')], 2.12 [dd, J = 17.5 and 4.0 Hz, CH₂H_bCH], 1.94 [dd, J = 17.5 and 11.0 Hz,

CH₂H_bCH], 1.38 [d, J = 7.0 Hz, CH₃(1)], 1.00 [d, J = 6.5 Hz, CH₃(3)]; ¹H NMR of 121b (500 MHz, CDCl₃): δ 7.39-7.21 [m, benzyl ArH], 7.12 [d, J = 8.0 Hz, ArH(7')], 7.06 [d, J = 8.0 Hz, ArH(6')], 6.79 [s, ArH(1')], 6.69 [s, ArH(3')], 6.48 [s, ArH(7)], 5.30 [s, OCH₂OCH₃], 4.01 [q, J = 6.5 Hz, NCHCH₃], 3.97 [s, O(4')CH₃], 3.84 [s, O(6)CH₃], 3.76 [d, J = 14.0 Hz, NCH₂H_bPh], 3.65 [s, O(8)CH₃], 3.62 [s, OCH₂OCH₃], 3.36 [ddq, J = 12.0, 6.5 and 4.5 Hz, CH₂H_bCHCH₃], 3.31 [d, J = 14.5 Hz, NCH₂H_bPh], 2.35 [s, CH₃(2')], 2.17 [dd, J = 18.0 and 12.0 Hz, CH₂H_bCH], 1.82 [dd, J = 17.5 and 4.5 Hz, CH₂CH_bCH], 1.38 [d, J = 7.0 Hz, CH₃(1)], 1.03 [d, J = 6.5 Hz, CH₃(3)].

Example 20. General Procedure for the Hydrolysis Reactions

To a stirred solution of methoxymethyl-protected starting material in a mixed solvent (MeOH/CH₂Cl₂-10:1, 0.01 M) was added 10 N HCl (1/20 volume of solvent). The reaction mixture was stirred at room temperature for 16 h. After this period of time the solvent was evaporated. EtOAc and saturated NaHCO₃ were added. The organic layer was washed with brine, dried over Na₂SO₄, and concentrated in vacuo. The crude product was purified by chromatography.

A. Preparation of Compound (15)




25

Compound 15 was obtained from compound 14 in 71 % yield. The product was purified by MPLC (hexanes/EtOAc/Et₃N; 3:1:0.1); ¹H NMR of S-15 (500 MHz, CDCl₃): δ 9.40 [s, OH], 7.39-6.95 [m, ArH(7') and benzyl ArH], 6.91

30

[d, $J = 8.0$ Hz, ArH(6')], 6.76 [s, ArH(1')], 6.62 [s, ArH(3')], 6.52 [s, ArH(7)], 5.01 [s, O(6)CH₂Ph], 4.88 [d, $J = 13.0$ Hz, O(8)CH₂H_bPh], 4.82 [d, $J = 13.0$ Hz, O(8)CH₂H_bPh], 4.08 [q, PhCHCH₃, hidden by O(4')CH₃], 4.08 [s, O(4')CH₃], 3.72 [d, $J = 14.0$ Hz, NCH₂H_bPh], 3.37 [ddq, $J = 11.5, 4.0$, and 6.5 Hz, CH_aH_bCHCH₃], 3.29 [d, $J = 14.0$ Hz, NCH₂H_bPh], 2.36 [s, CH₃(2')], 2.21 [dd, $J = 17.5$ and 4.0 Hz, CH₂H_bCH], 1.90 [dd, $J = 17.5$ and 11.5 Hz, CH₂H_bCH], 1.40 [d, $J = 6.5$ Hz, CH₃(1)], and 1.01 [d, $J = 6.5$ Hz, CH₃(3)]; ¹H NMR of R-15 (500 MHz, CDCl₃): δ 9.42 [s, OH], 7.39-6.95 [m, ArH(7') and benzyl ArH], 6.90 [d, $J = 8.0$ Hz, ArH(6')], 6.85 [s, ArH(1')], 6.63 [s, ArH(3')], 6.50 [s, ArH(7)], 5.03 [d, $J = 11.5$ Hz, O(6)CH₂H_bPh], 4.97 [d, $J = 11.5$ Hz, O(6)CH₂H_bPh], 4.87 [d, $J = 13.0$ Hz, O(8)CH₂H_bPh], 4.82 [d, $J = 13.0$ Hz, O(8)CH₂H_bPh], 4.08 [q, PhCHCH₃, hidden by O(4')CH₃], 4.08 [s, O(4')CH₃], 3.77 [d, $J = 14.0$ Hz, NCH₂H_bPh], 3.37 [ddq, $J = 11.5, 4.0$, and 6.5 Hz, CH_aH_bCHCH₃], 3.34 [d, $J = 14.0$ Hz, NCH₂H_bPh], 2.36 [s, CH₃(2')], 2.24 [dd, $J = 17.5$ and 11.5 Hz, CH₂H_bCH], 1.90 [dd, $J = 17.5$ and 4.0 Hz, CH₂H_bCH], 1.38 [d, $J = 6.5$ Hz, CH₃(1)], and 1.05 [d, $J = 6.5$ Hz, CH₃(3)].

B. Preparation of Compound (122)



In a 10 mL round bottom flask, a mixture of substrates **121a** and **121b** (11 mg, 0.02 mmol) was dissolved in methanol (3 mL). To the solution was added 10% aqueous HCl (2 mL) and the reaction mixture was stirred at room temperature for 12 h. The reaction mixture was concentrated in vacuo and the resulting residue was dissolved in ethyl acetate (10 mL), washed with sodium

bicarbonate, water, and dried over sodium sulfate. Concentration in vacuo yielded a mixture of compounds **122a** and **122b** (10 mg, 100%) as a white solid.


Example 21. General Procedure for the Per-debenzylation Reactions

5

To a solution of benzyl-protected monomer in a mixed solvent (MeOH/CH₂Cl₂-2:1, 0.01 M) was added 10 % Pd/C (20 mol %). The atmosphere was exchanged for N₂, then H₂, and then a H₂ balloon was attached. The reaction mixture was stirred until TLC analysis indicated no starting material and possible
 10 intermediate left. The catalyst was removed by passing through a bed of Celite. The filtrate was concentrated in vacuo to yield deprotected monomer. The mixture of atropisomers was able to be separated by HPLC using an amino-bonded column.

A. Preparation of Korupensamine A & "C"

15



20

Compound **4** and **4'** were obtained from compounds **15** in 75 % yield. The atropisomers Korupensamine A [**4**] and "Korupensamine C" [**4'**] were separated by HPLC using an amino-bonded column (CHCl₃/MeOH/(NH₄)₂CO₃; 95:5:0.1); ¹H NMR of **4** (HOAc Salt) (500 MHz, CD₃OD, referenced to CHD₂OD @ 3.30 ppm): δ 7.09 [d, J = 8.0 Hz, ArH(7')], 6.80 [d, J = 8.0 Hz, ArH(6')], 6.78 [s, ArH(3')], 6.69 [s, Ar(1')], 6.44 [s, ArH(7)], 4.75 [q, J = 7.0 Hz, ArCHCH₃], 4.08 [s, O(4')CH₃], 3.65 [ddq, J = 12.0, 5.0 and 6.5 Hz, CH₂H_bCHCH₃], 2.62 [dd, J = 18.0 and 5.0 Hz, CH_aH_bCH], 2.30 [s, CH₃(2')], 2.05 [dd, J = 18.0 and 12.0 Hz, CH_aH_bCH], 1.64 [d, J = 7.0 Hz, CH₃(1)], and 1.19 [d, J = 6.5 Hz, CH₃(3)]; ¹H NMR of **4'** (HOAc Salt) (500 MHz, CD₃OD, referenced to CHD₂OD @ 3.30 ppm): δ 7.02 [d, J = 8.0 Hz, ArH(7')], 6.80 [d, J = 8.0 Hz, ArH(6')], 6.80 [s, ArH(1' or 3')], 6.78 [s, Ar(3' or 1')], 6.44 [s, ArH(7)], 4.74 [q, J = 7.0 Hz,

30

ArCHCH₃], 4.08 [s, O(4')CH₃], 3.62 [ddq, J = 12.0, 5.0 and 6.5 Hz, CH₂H_bCHCH₃], 2.38 [dd, J = 18.0 and 12.0 Hz, CH₂H_bCH], 2.33 [s, CH₃(2')], 2.23 [dd, J = 18.0 and 5.0 Hz, CH₂H_bCH], 1.67 [d, J = 6.5 Hz, CH₃(1)], and 1.23 [d, J = 6.5 Hz, CH₃(3)].

5

Example 22. General Procedure for the Silver Oxide Promoted Oxidative Coupling and Simultaneous Reductive Bleaching/Perdebenzylation Reactions


10

15

20


To a stirred solution of benzyl-protected monomer in CH₂Cl₂ (0.01 M) was added 5 equivalent of Ag₂O. The reaction mixture was stirred at room temperature in the dark for 40 h. The solid was removed by passing through the Celite bed. MeOH (volume equal to that of CH₂Cl₂) was added to the filtrate, followed by the addition of 10 % Pd/C (20 mol %). The atmosphere was exchanged for N₂, then H₂, and then a H₂ balloon was attached. The reaction mixture was stirred until TLC analysis indicated no starting material and possible intermediate left. The catalyst was removed by passing through a bed of Celite. The filtrate was concentrated to yield deprotected dimer. The crude product was further purified by HPLC with amino-bonded column.

A. Preparation of Michellamines A, B, and C



- Michellamines A-C [1-3] were obtained from compound 15 in 90 % yield. They were separated by HPLC using an amino-bonded column
- 5 (CHCl₃/MeOH/(NH₄)₂CO₃; 93:7:0.1); ¹H NMR of Michellamine A (1, HOAc Salt) (500 MHz, CD₃OD, referenced to CHD₂OD @ 3.30 ppm): δ 7.30 [s, ArH(7')], 6.85 [s, ArH(3')], 6.74 [s, ArH(1')], 6.44 [s, ArH(7)], 4.77 [q, J = 7.0 Hz, ArCHCH₃], 4.10 [s, O(4')CH₃], 3.70 [ddq, J = 12.0, 4.5 and 6.5 Hz, CH_aH_bCHCH₃], 2.82 [dd, J = 18.0 and 4.5 Hz, CH_aH_bCH], 2.34 [s, CH₃(2')], 2.15 [dd, J = 18.0 and 12.0 Hz, CH_aH_bCH], 1.65 [d, J = 6.5 Hz, CH₃(1)], and 1.24 [d, J = 6.5 Hz, CH₃(3)]; ¹H NMR of Michellamine B (2, HOAc Salt) (500 MHz, CD₃OD, referenced to CHD₂OD @ 3.30 ppm): δ 7.32/7.27 [s, ArH(7')], 6.86/6.74 [s, ArH(3')], 6.85/6.83 [s, ArH(1')], 6.45 [s, ArH(7)], 4.76/4.73 [q, J = 7.0/7.0 Hz, ArCHCH₃], 4.10/4.09 [s, O(4')CH₃], 3.73-3.62 [m, CH_aH_bCHCH₃], 2.79 [dd, J = 17.5 and 5.0 Hz, CH_aH_bCH], 2.53 [dd, J = 18.0 and 11.5 Hz, CH_aH_bCH], 2.36/2.33 [s, CH₃(2')], 2.34-2.29 [dd, CH_aH_bCH, hidden by CH₃(2')], 2.12 [dd, J = 18.0 and 11.5 Hz, CH_aH_bCH], 1.69/1.64 [d, J = 6.5/7.0 Hz, CH₃(1)], and 1.26/1.22 [d, J = 6.0/6.5 Hz, CH₃(3)].
- 10
- 15

B. Preparation of Compounds (123a), (123b), and (123c)



10

In a 10 mL round bottom flask, substrates 122a and 122b (5 mg, 0.01mmol) were dissolved in methylene chloride (3 mL). To the solution was added silver (I) oxide (7 mg, 0.03 mmol) and the mixture was stirred at room temperature overnight. When TLC analysis showed no starting materials were left, the mixture was filtered through a bed of Celite bed. Concentration of the filtrate gave a blue solid (5 mg, 100%) which was dissolved in a mixed solvent of methylene chloride (2 mL) and methanol (2 mL). To the mixture was added a solution of NaBH₄ in methanol (2 mL). The mixture was concentrated, and the residue was dissolved in methylene chloride (10 mL), washed with water, and dried over sodium sulfate.

15


20

Concentration in vacuo yielded a mixture of compounds 123a, 123b, and 123c (5 mg, 100%) as a white solid.

5 It will be appreciated by those skilled in the art that various modifications can be made to the above described embodiments of the invention without departing from the essential nature thereof. The invention is intended to encompass all such modifications within the scope of the appended claims.

33

1. A compound of the formula (I):



10 wherein X is Br, Cl or I, Y is H, (C₁-C₄)alkyl, benzyl, or CHO, each of R¹ and R² is H or CH₃, R³ is H or (C₂-C₅)acyl and R⁴ is a protecting group selected from the group consisting of (C₁-C₄)alkyl, benzyl, (C₂-C₅)acyl and an acid-labile hydroxy protecting group.

- 15 2. The compound of claim 1 wherein R⁴ is an acid-labile hydroxy protecting group.

3. The compound of claim 2 wherein the acid labile protecting group is (C₁-C₄)alkoxy(C₁-C₄)alkyl, tetrahydropyranyl, or (R⁸)₃Si, wherein each R⁸ is (C₁-C₄)alkyl.

- 20 4. The compound of claim 1 wherein R¹ and R² are CH₃.


5. The compound of claim 4 wherein C₁ and C₃ have the *R* configuration.

6. The compound of claim 4 wherein X=I.

25

7. A compound of the formula (I):

34



(I)

wherein X is Br, Cl or I, Y is H, (C₁-C₄)alkyl, benzyl, or CHO, each of R¹ and R² is H or CH₃, R³ is a protecting group selected from the group consisting of (C₁-C₄)alkyl, benzyl, (C₂-C₅)acyl and an acid-labile hydroxy protecting group; and R⁴ is H or (C₂-C₅)acyl.

8. The compound of claim 7 wherein R³ is an acid-labile hydroxy protecting group.

9. The compound of claim 8 wherein the acid labile protecting group is (C₁-C₄)alkoxy(C₁-C₄)alkyl, tetrahydropyranyl, or (R⁸)₃Si, wherein each R⁸ is (C₁-C₄)alkyl.


10. The compound of claim 7 wherein R¹ and R² are CH₃.

11. The compound of claim 10 wherein C₁ and C₃ have the *R* configuration.

12. The compound of claim 10 wherein X=I.

13. A compound of the formula (II):

35




(II)

wherein R^6 is Cl, Br, I, $B(OH)_2$, an anhydride or ester of $B(OH)_2$, or OSO_2R^9 , wherein R^9 is (C_1-C_4) perfluoroalkyl, and each of R^5 and R^7 is H, (C_1-C_4) alkyl, benzyl, (C_2-C_5) acyl or an acid-labile hydroxy protecting group.

14. The compound of claim 13 wherein R^6 is Br or $B(OH)_2$, R^5 is an acid-labile protecting group, and R^7 is H or CH_3 .


15. A method of preparing a korupensamine or an analog thereof comprising:
(a) reacting a compound of the formula (III):



(III)


wherein each of R^1 and R^2 is CH_3 or H, X is I, Y is (C_1-C_4) alkyl, benzyl or CHO, and each of R^3 and R^4 is (C_1-C_4) alkyl, benzyl, (C_2-C_5) acyl or an acid-labile hydroxy protecting group; with a compound of the formula (IV):

36



(IV)

wherein R^5 is benzyl, (C_2-C_5) acyl or an acid-labile hydroxy protecting group, R^6 is $B(OH)_2$, and R^7 is (C_1-C_4) alkyl; in the presence of a Pd(0) catalyst and an inorganic base in an organic solvent, to yield a compound of the formula (V):



(V)

wherein Y, R^1 , R^2 , R^3 , R^4 , R^5 and R^7 are as defined above for compounds of formula III and IV.

16. The method of claim 15 further comprising (b) removing protecting groups R^3 , R^4 , R^5 and Y to yield a compound of formula V wherein each of R^1 and R^2 is H or CH_3 , R^7 is (C_1-C_4) alkyl, and Y, R^2 , R^3 , R^4 , and R^5 are H.

17. The method of claim 16 wherein C_1 and C_3 have the *R* configuration.

18. The method of claim 17 wherein the korupensamine prepared is korupensamine A or korupensamine B.
19. The method of claim 15 wherein R^1 , R^2 and R^7 are CH_3 , R^5 is an acid-labile hydroxy protecting group that is subsequently removed by exposing V to dilute aqueous acid, and Y, R^3 and R^4 are benzyl that are subsequently removed by hydrogenolysis.
20. The method of claim 19 wherein R^5 is methoxymethyl.
21. The method of claim 15 wherein C_1 and C_3 have the *R* configuration, R^1 , R^2 and R^7 are CH_3 , Y, R^3 and R^4 are benzyl, and R^5 is an acid-labile hydroxy protecting group, further comprising (b) removing the acid-labile hydroxy protecting group by exposing V to dilute acid and (c) oxidatively coupling two molecules of V to yield *S,S*-16, *R,S*-16 and *R,R*-16 as shown in Fig. 1.
22. The method of claim 21 further comprising (d) reducing the compounds of formula 16 and removing the benzyl groups to yield at least one michellamine.




FIG. 1

INTERNATIONAL SEARCH REPORT

Intern. Application No
PCT/US 95/14896

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D217/24 C07C39/38 C07C309/66 C07F5/02 C07F5/04
C07F5/05

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JOURNAL OF THE CHEMICAL SOCIETY, no. 4, 1963 pages 3940-3945, XP 000561811 HORII Z. ET AL. '748. The synthesis of musizin' see page 3942, formula XVIII ---	13,14
P,X	TETRAHEDRON LETTERS, vol. 35, no. 47, 21 November 1994 pages 8747-8750, XP 000562381 HOYE T.R. ET AL. 'Total synthesis of michellamines A-C: Important anti-HIV agents' see page 8748, compounds 12 and 13	13,14
A	see page 8748, compound 8	1-12
A	see scheme on page 8749 ---	15-22
	-/-	

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

11 March 1996

Date of mailing of the international search report

3.04.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hartrampf, G

INTERNATIONAL SEARCH REPORT

 Inter. Application No
 PCT/US 95/14896

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	HETEROCYCLES, vol. 39, no. 2, 31 December 1994 pages 503-508, XP 000561812 BRINGMANN G. ET AL. 'First total synthesis of korupensamines A and B' see page 505, compound 7	13,14
A	see page 505, compound 10	1-12
A	see scheme on page 506	15-22
A	--- ANGEWANDTE CHEMIE, INTERNATIONAL EDITION IN ENGLISH, vol. 25, no. 10, 1986 pages 913-915, BRINGMANN G. ET AL. 'Regioselective and atropisomeric-selective aryl coupling to give naphthyl iosquinoline alkaloids: The first total synthesis of (-)-ancistrocladine' cited in the application see the whole document	1-12
A	--- AUSTRALIAN JOURNAL OF CHEMISTRY, vol. 43, no. 1, 1990 pages 79-86, XP 000562132 RIZZACASA M.A. ET AL. 'The stereoisomers of 5-bromo-5,8-dimethoxy-1,2,3-trimethyl-1,2, 3,4-tetrahydroisoquinoline: X-ray crystal structure of the trans isomer' see page 80, compounds 18 and 20	1-12
A	--- LIEBIGS ANNALEN DER CHEMIE, 1993 pages 877-888, BRINGMANN G. ET AL. 'The synthesis of all possible isomeric 6,8-dioxygenated 1,3-dimethyl-1,2,3,4-tetrahydroisoquinolin e methyl ethers - useful chiral building blocks for naphthyl iosquinoline alkaloids' cited in the application see the whole document -----	1-12